



## Fracture Liaison Service Database

Annual report 12 January 2023

### References and bibliography

Ben-Shlomo Y, Bhimjiyani A, Gregson CL, Patel R. Social deprivation predicts adverse health outcomes after hospital admission with hip fracture in England. *Osteoporosis Int.* 2021;32:1129-1141.

Brennan-Olsen SL, Buchbinder R, SM Hosking, JA Pasco, LJ Williams. The Role of Health Literacy in the Treatment of Osteoporosis. *J Bone Miner Res.* 2016;31:1909.

International Osteoporosis Foundation. Broken bones, broken lives: A roadmap to solve the fragility fracture crisis in Europe. Nyon: IOF, 2018.

National Institute for Health and Care Excellence. Bisphosphonates for treating osteoporosis. NICE technology appraisal guidance (TA464). Manchester: NICE, 2017. [www.nice.org.uk/guidance/ta464](http://www.nice.org.uk/guidance/ta464)

National Institute for Health and Care Excellence. Denosumab for the prevention of osteoporotic fractures in postmenopausal women. NICE technology appraisal guidance (TA204). Manchester: NICE, 2010. [www.nice.org.uk/guidance/ta204](http://www.nice.org.uk/guidance/ta204)

National Institute for Health and Care Excellence. Falls in older people: assessing risk and prevention. NICE clinical guideline (CG161). Manchester: NICE, 2013. [www.nice.org.uk/guidance/cg161](http://www.nice.org.uk/guidance/cg161)

National Institute for Health and Care Excellence. Falls in older people. NICE clinical guideline (QS86). Manchester: NICE, 2015. <https://www.nice.org.uk/guidance/qs86>

National Institute for Health and Care Excellence. Osteoporosis: assessing the risk of fragility fracture. NICE clinical guideline (CG146). Manchester: NICE, 2012. <https://www.nice.org.uk/guidance/cg146>

National Institute for Health and Care Excellence. Osteoporosis. NICE quality standard (QS149). Manchester: NICE, 2017. [www.nice.org.uk/guidance/qs149](http://www.nice.org.uk/guidance/qs149)

National Institute for Health and Care Excellence. Raloxifene and teriparatide for the secondary prevention of osteoporotic fragility fractures in postmenopausal women. NICE clinical guideline (TA161). Manchester: NICE, 2008. <https://www.nice.org.uk/guidance/ta161>

National Institute for Health and Care Excellence. Romosozumab for treating severe osteoporosis. NICE technology appraisal guidance (TA791). Manchester: NICE, 2022. <https://www.nice.org.uk/guidance/ta791>

National Osteoporosis Guideline Group. NOGG 2021: Clinical guideline for the prevention and treatment of osteoporosis. Sheffield: NOGG, 2021. [www.sheffield.ac.uk/NOGG/](http://www.sheffield.ac.uk/NOGG/)



Royal College  
of Physicians

FLS-DB

Royal College of Physicians  
11 St Andrews Place, Regent's Park  
London NW1 4LE

+44 (0)20 3075 1511  
flsdb@rcp.ac.uk  
[rcp.ac.uk](http://rcp.ac.uk)

Royal College of Physicians. Fracture Liaison Service Database annual report. Benchmarking FLS improvement and performance in 2019: Pre-COVID. London: RCP, 2021.

Royal Osteoporosis Society. Effective secondary prevention of fragility fractures: clinical standards for fracture liaison services. Bath: ROS, 2019. <https://theros.org.uk/media/1eubz33w/ros-clinical-standards-for-fracture-liaison-services-august-2019.pdf>

Royal Osteoporosis Society. FLS Implementation Toolkit. Bath: ROS, 2022.  
<https://theros.org.uk/healthcare-professionals/fracture-liaison-services/implementation-toolkit/>.

Royal Osteoporosis Society. Quality Standards for Osteoporosis and Prevention of Fragility Fractures. Bath: ROS, 2017. <https://theros.org.uk/media/99099/op-standards.pdf>